



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/26	A1	(11) International Publication Number: WO 99/29336 (43) International Publication Date: 17 June 1999 (17.06.99)
<p>(21) International Application Number: PCT/US98/25515</p> <p>(22) International Filing Date: 2 December 1998 (02.12.98)</p> <p>(30) Priority Data: 60/067,600 5 December 1997 (05.12.97) US</p> <p>(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): HOFFMANN, James, Arthur [US/US]; 4272 Woodland Streams Drive, Greenwood, IN 46143 (US).</p> <p>(74) Agents: MACIAK, Ronald, S. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: GLP-1 FORMULATIONS</p> <p>(57) Abstract</p> <p>Methods and formulations are presented that provide for a) the oral absorption of GLP-1 peptides that bind surfactants; and b) long-term storage of formulations containing these peptides. For example, a GLP-1/DSS complex is administered orally instead of parenterally, which is much more convenient for, and facilitates compliance with diabetic patients and persons with other GLP-1 treated conditions.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

GLP-1 FORMULATIONS

BACKGROUND OF THE INVENTION

Formulations are presented that have improved storage characteristics. These formulations are particularly
5 suitable for oral absorption of GLP-1 peptides that bind surfactants.

Administration of therapeutic peptides is often limited to parenteral routes rather than preferred oral administration due, e.g. to destruction of the peptides if
10 ingested rather than injected. This is unfortunate because many peptides have proven clinically effective and could have more widespread use if easy to administer and acceptable to recipients. For example, GLP-1-like molecules possess anti-diabetic activity in human subjects suffering
15 from Type II and, in some cases, even Type I diabetes. Treatment with GLP-1 elicits activity (increased insulin secretion and biosynthesis, reduced glucagon secretion, delayed gastric emptying) only at elevated glucose levels, and thus provides a potentially much safer therapy than
20 insulin or sulfonylureas. Post-prandial and glucose levels in patients can be moved toward normal levels with proper GLP-1 therapy. There are also reports suggesting GLP-1-like molecules possess the ability to preserve and even restore pancreatic beta cell function in Type-II patients. On the
25 other hand, to be effective as a treatment, GLP-1 formulations may have to be administered by injection at, or slightly before, each meal. This is the regimen used to administer insulin. For such a regimen, a multi-use solution formulation stored for long periods of time at

-2-

refrigerated or ambient temperature is preferred. Such a formulation must contain a preservative with sufficient anti-microbial properties to prevent degradation and contamination of the solution. Unfortunately, preservatives
5 tend to deleteriously affect the therapeutic agent, e.g. a peptide. For example, solutions of GLP-1 molecules undergo conformational changes in the presence of a preservative such as phenol. In the presence of the preservative meta-cresol (m-cresol), aqueous solutions of GLP-1 molecules that
10 are near neutral pH turn hazy, and precipitation develops. What is needed therefore, are additives for formulations of peptides such as GLP-1 molecules that allow storage at refrigeration (about 4°C or lower) and/or ambient temperatures while still preserving both solution clarity,
15 compound integrity, and biological activity.

SUMMARY OF THE INVENTION

Methods and formulations of the present invention provide formulations for a. oral absorption of GLP-1 peptides that bind surfactants with high affinity; b. long
20 term storage of formulations containing these peptides.

An aspect of the invention is a formulation comprising a GLP-1 peptide and a small quantity of a surfactant. Preferred surfactants include DSS (docusate sodium, CAS Registry Number [577-11-7]) and related substances; docusate
25 calcium [CAS number 128-49-4], and docusate potassium [CAS number 7491-09-0]. Other surfactants include SDS (sodium dodecyl sulfate or sodium lauryl sulfate), sodium caprylate, sodium cholate, sodium deoxycholate, sodium taurocholate,

-3-

and sodium glycocholate. Suitable agents also include zwitterionic (e.g. N-alkyl-N,N-dimethylammonio-1-propanesulfonates, 3-cholamido-1-propyldimethylammonio-1-propane-sulfonate), cationic (cetylpyridinium chloride),
5 non-ionic (Triton X-100, Dodecyl β -D-glucopyranoside), or polymeric (Tween-40, Tween-80, Brij-35) surfactants.

Peptides used in the formulations of the present invention include GLP-1 or GLP-1-like molecules. A preferred GLP-1-like molecule is Val⁸-GLP-1. Other suitable
10 GLP-1-like molecules include the 2 native GLP-1 forms, position-8 analogs, and molecules containing a C-terminal acid.

The formulation is stable at a pH of about 6.5 to 9.0, more preferably at a pH of about 7 to 8. The formulation
15 includes a preservative. Preferred preservatives include m-cresol, phenol, methylparaben, and benzyl alcohol. The formulation is stable during long term storage at 4°C and ambient temperature. The formulation optionally includes an isotonicity agent, for example glycerin, or sodium chloride.

20 Another aspect of the invention is a method of treating a person having diabetes or other conditions in which the administration of a GLP-1-like molecule is indicated. The method includes obtaining a formulation of the present invention and administering a pharmacologically effective
25 amount of the formulation to the person. Preferably an oral route is used to administer the formulation, although a parenteral route is also suitable.

DETAILED DESCRIPTION OF THE INVENTION

-4-

Methods and formulations of the present invention provide for a) the oral absorption of GLP-1 peptides that bind surfactants with high affinity; and b) long-term storage of preserved formulations containing these peptides.

5 In an embodiment of the invention, a GLP-1/DSS complex is used to administer GLP-1 orally instead of parenterally. This aspect of the invention provides much greater convenience and compliance for diabetic patients and persons having other conditions in which treatment with a GLP-1-like
10 molecule is indicated. This characteristic will make GLP-1 treatment more useful and widely available. Use of preservatives prevents microbial contamination and therefore allows multiple use from a single solution.

Several key observations suggest that a significant
15 portion of a GLP-1 peptide in a formulation containing sodium docusate (DSS) will be absorbed orally:

- a. DSS binds to GLP-1 with a high affinity;
- b. DSS binding alters GLP-1 secondary structure; this altered structure may correspond to a membrane-transportable
20 state as described by Milstein (1996). The DSS appears to be acting as a so-called carrier molecule.
- c. After administration of the formulation into a body (subcutaneously) the GLP-1 peptide exhibits full biological activity; this suggests either that the GLP-1 in
25 the formulation retains its receptor binding affinity or the GLP-1-DSS complex in the formulation can be disrupted, reforming the native GLP-1 in an alpha-helix structure; a CD study showed that a 2-day dialysis of a GLP-1-DSS mixture did not revert the GLP-1 back to its alpha-helix
30 conformation.

-5-

d. Large quantities of DSS can be safely administered orally because it is already approved for use as a laxative in humans; some of the orally administered DSS is absorbed systemically.

5 The addition of an anionic surfactant sodium docusate (DSS), at a very low level (2:1 on a molar basis vs. peptide), also dramatically improved the solution stability of Val⁸-GLP-1(7-37)OH in a formulation that is isotonic, is at a near neutral pH (pH 7.8), and also contains a suitable
10 preservative (m-cresol). This formulation provides an improved product that should meet antimicrobial-sterility standards throughout the world. Improvement in formulation stability is over a wide range of storage conditions, from about 2°C to about 37°C, more preferably at about 4° to
15 about 25°C.

 In an embodiment, the formulation allows single or multi-use parenteral formulation of a GLP-1 analog to be prepared that is suitable for long-term storage. Also, because the DSS facilitates the GLP-1 existing in a soluble
20 micelle or aggregated state, this formulation provides an improved prolonged time action after subcutaneous administration.

 The anionic surfactant, sodium docusate (DSS) has a very high affinity for a GLP-1 compound, specifically
25 Val⁸-GLP-1(7-37)OH and, upon binding to the peptide, the Val⁸-GLP-1 secondary structure is converted from mostly alpha-helix to mostly a beta sheet. A slightly larger form of Val⁸-GLP-1 with DSS molecule(s) bound to it was observed on size-exclusion chromatography (SEC) and the altered

-6-

secondary structure was noted by circular dichroism experiments (CD).

A formulation containing DSS and Val⁸-GLP-1 injected subcutaneously into dogs showed insulintropic-like activity comparable in potency to Val⁸-GLP-1 in a phosphate buffer solution (PBS formulation).

Preferred embodiments for a surfactant include DSS (docusate sodium, CAS Registry Number [577-11-7]) and related substances; docusate calcium [CAS number 128-49-4], docusate potassium [CAS number 7491-09-0].

Also preferred are other surfactants including: SDS (sodium dodecyl sulfate or sodium lauryl sulfate), sodium caprylate, sodium cholate, sodium deoxycholate, sodium taurocholate, and sodium glycocholate.

Other suitable surfactants include: zwitterionic (e.g. N-alkyl-N,N-dimethylammonio-1-propanesulfonates, 3-cholamido-1-propyldimethylammonio-1-propane-sulfonate), cationic (cetylpyridinium chloride), non-ionic (Triton X-100, Dodecyl β -D-glucopyranoside), or polymeric (Tween-40, Tween-80, Brij-35) surfactants.

Preferred preservatives include m-cresol and phenol. Also preferred are methylparaben, benzyl alcohol, and other similar preservatives.

A preferred isotonicity agent is glycerin, also preferred is any isotonicity agent (e.g. sodium chloride).

Optionally, a wide range of excipients may be included in the formulation, such as glycerin, m-cresol, phenol, methylparaben, and the like, although the excipients alone would not provide the dramatic improvement in solution stability that characterizes the present invention. Some of

-7-

these excipients are preservatives, some are isotonicity agents.

GLP-1-like molecules include GLP-1 analogs and derivatives, GLP-1 molecules, native as well as GLP-1
5 analogs, that bind tightly (that is, with high affinity) with surfactants. A preferred GLP-1 molecule is: Val⁸-GLP-1.

GLP-1 molecules such as native GLP-1(7-36)NH₂ and GLP-1(7-37)OH, as well as other GLP-1 analogs are also suitable
10 for the practice of the invention. Also preferred are position-8 analogs and analogs containing a C-terminal acid. All other analogs are also suitable if they bind with high affinity to surfactants.

"GLP-1" means GLP-1(7-37). By custom in the art, the
15 amino-terminus of GLP-1(7-37) has been assigned number 7 and the carboxy-terminus has been assigned number 37. The amino acid sequence of GLP-1(7-37) is well-known in the art, but is presented as SEQ ID NO:1 for the reader's convenience.

A "GLP-1 analog" is defined as a molecule having one or
20 more amino acid substitutions, deletions, inversions, or additions compared with GLP-1. GLP-1 analogs known in the art include, for example, GLP-1(7-34), GLP-1(7-35), GLP-1(7-36), Val⁸-GLP-1(7-37), Gln⁹-GLP-1(7-37), D-Gln⁹-GLP-1(7-37), Thr¹⁶-Lys¹⁸-GLP-1(7-37), and
25 Lys¹⁸-GLP-1(7-37).

A "GLP-1 derivative" is defined as a molecule having the amino acid sequence of GLP-1 or of a GLP-1 analog, but additionally having chemical modification of one or more of its amino acid side groups, α -carbon atoms, terminal amino
30 group, or terminal carboxylic acid group. A chemical

-8-

modification includes, but is not limited to, adding chemical moieties, creating new bonds, and removing chemical moieties. Modifications at amino acid side groups include, without limitation, acylation of lysine ϵ -amino groups, N-alkylation of arginine, histidine, or lysine, alkylation of glutamic or aspartic carboxylic acid groups, and deamidation of glutamine or asparagine. Modifications of the terminal amino include, without limitation, the des-amino, N-lower alkyl, N-di-lower alkyl, and N-acyl modifications. Modifications of the terminal carboxy group include, without limitation, the amide, lower alkyl amide, dialkyl amide, and lower alkyl ester modifications. Lower alkyl is C(1)-C(4) alkyl. Furthermore, one or more side groups, or terminal groups, may be protected by protective groups known to the ordinarily-skilled protein chemist. The α -carbon of an amino acid may be mono- or di-methylated.

The use in the present invention of a molecule claimed in U.S. Patent No. 5,120,712, GLP-1(7-37)OH, which is expressly incorporated by reference, is highly preferred. Such molecule is selected from the group consisting of a peptide having the amino acid sequence of SEQ ID NO: 1 and a derivative of said peptide, wherein said peptide is selected from the group consisting of: a pharmaceutically-acceptable acid addition salt of said peptide; a pharmaceutically-acceptable carboxylate salt of said peptide; a pharmaceutically-acceptable lower alkylester of said peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

-9-

A preferred group of GLP-1 analogs and derivatives for use in the present invention is composed of the various GLP-1 molecules claimed in U.S. Patent No. 5,545,618, which is herein expressly incorporated by reference.

5 A preferred group of GLP-1 analogs and derivatives for use in the present invention is composed of molecules of formula:

R_1 -SEQ ID NO:2- R_2

and the pharmaceutically-acceptable salts thereof, wherein:

10 R_1 is selected from the group consisting of L-histidine, D-histidine, desamino-histidine, 2-amino-histidine, β -hydroxy-histidine, homohistidine, alpha-fluoromethyl-histidine, and alpha-methyl-histidine; and R_2 is selected from the group consisting of NH_2 , and
15 Gly-OH.

Numerous such GLP-1 analogs and derivatives have been disclosed and include, for example:

GLP-1(7-36) NH_2 , Gly⁸-GLP-1(7-36) NH_2 , Gln⁹-GLP-1(7-37),
D-Gln⁹-GLP-1(7-37), acetyl-Lys⁹-GLP-1(7-37),
20 Thr⁹-GLP-1(7-37), D-Thr⁹-GLP-1(7-37), Asn⁹-GLP-1(7-37),
D-Asn⁹-GLP-1(7-37), Ser²²-Arg²³-Arg²⁴-Gln²⁶-GLP-1(7-37),
Thr¹⁶-Lys¹⁸-GLP-1(7-37), Lys¹⁸-GLP-1(7-37), Arg²³-GLP-1(7-37),
Arg²⁴-GLP-1(7-37), and the like (see, e.g., WO 91/11457).

Another preferred group of active compounds for use in
25 the present invention is disclosed in WO 91/11457, and consists essentially of GLP-1(7-34), GLP-1(7-35), GLP-1(7-36), or GLP-1(7-37), or the amide form thereof, and pharmaceutically-acceptable salts thereof, having at least one modification selected from the group consisting of:

-10-

- (a) substitution of at least one of the following
glycine, serine, cysteine, threonine, asparagine, glutamine,
tyrosine, alanine, valine, isoleucine, leucine, methionine,
phenylalanine, arginine, or D-lysine for lysine at position
5 26 and/or position 34; or substitution of glycine, serine,
cysteine, threonine, asparagine, glutamine, tyrosine,
alanine, valine, isoleucine, leucine, methionine,
phenylalanine, lysine, or a D-arginine for arginine at
position 36;
- 10 (b) substitution of an oxidation-resistant amino acid
for tryptophan at position 31;
- (c) substitution of at least one of the following:
tyrosine for valine at position 16; lysine for serine at
position 18; aspartic acid for glutamic acid at position
15 21; serine for glycine at position 22; arginine for
glutamine at position 23; arginine for alanine at position
24; and glutamine for lysine at position 26; and
- (d) substitution of at least one of the following:
glycine, serine, or cysteine for alanine at position 8;
20 aspartic acid, glycine, serine, cysteine, threonine,
asparagine, glutamine, tyrosine, alanine, valine,
isoleucine, leucine, methionine, or phenylalanine for
glutamic acid at position 9; serine, cysteine, threonine,
asparagine, glutamine, tyrosine, alanine, valine,
25 isoleucine, leucine, methionine, or phenylalanine for
glycine at position 10; and glutamic acid for aspartic acid
at position 15; and
- (e) substitution of at least one of the following:
glycine, serine, cysteine, threonine, asparagine, glutamine,
30 tyrosine, alanine, valine, isoleucine, leucine, methionine,

-11-

or phenylalanine, or the D- or N-acylated or alkylated form of histidine for histidine at position 7; wherein, in the substitutions is (a), (b), (d), and (e), the substituted amino acids can optionally be in the D-form and the amino acids substituted at position 7 can optionally be in the N-acylated or N-alkylated form.

Because the enzyme, dipeptidyl-peptidase IV (DPP IV), may be responsible for the observed rapid in vivo inactivation of administered GLP-1, (Mentlein et al. 1993), administration of GLP-1 analogs and derivatives that are protected from the activity of DPP IV is preferred, and the administration of Gly⁸-GLP-1(7-36)NH₂, Val⁸-GLP-1(7-37)OH, α-methyl-Ala⁸-GLP-1(7-36)NH₂, and Gly⁸-Gln²¹-GLP-1(7-37)OH, or pharmaceutically-acceptable salts thereof, is more preferred.

Another preferred group of molecules for use in the present invention consists of compounds, claimed in U.S. Patent No. 5,512,549, which is expressly incorporated herein by reference. This group is defined by the general formula:



and pharmaceutically-acceptable salts thereof, wherein R₁ is selected from the group consisting of 4-imidazopropionyl, 4-imidazoacetyl, or 4-imidazo-α, α dimethyl-acetyl; R₂ is selected from the group consisting of Gly-OH or NH₂. In addition, Lys at position 27 of SEQ ID NO:3 may be an acyl group selected from the group consisting of C₆-C₁₀ unbranched acyl or may be absent.

-12-

More preferred compounds of SEQ ID NO:3 for use in the present invention are those in which Xaa is Arg and Lys at position 27 is C₆-C₁₀ unbranched acyl.

Highly preferred compounds of SEQ ID NO:3 for use in the present invention are those in which Xaa is Arg, Lys at position 27 is C₆-C₁₀ unbranched acyl, and R₂ is Gly-OH.

More highly preferred compounds of SEQ ID NO:3 for use in the present invention are those in which Xaa is Arg, Lys at position 27 is C₆-C₁₀ unbranched acyl, R₂ is Gly-OH, and R¹ is 4-imidazopropionyl.

The most preferred compound of SEQ ID NO:3 for use in the present invention is that in which Xaa is Arg, Lys at position 27, is C⁸ unbranched acyl, R₂ is Gly-OH, and R¹ is 4-imidazopropionyl.

The use of GLP-1(7-36) amide, SEQ ID NO: 4, or a pharmaceutically-acceptable salt thereof, in the present invention is also highly preferred. The use of Val⁸-GLP-1(7-37)OH, SEQ ID NO:5, or a pharmaceutically-acceptable salt thereof, in the present invention is most highly preferred.

Other non-GLP-1 related peptides that bind DSS may also be made orally absorbable by the methods and formulations of the present invention. To determine whether these peptides are candidates for the formulations presented herein, it is useful to determine whether they bind with high affinity to a surfactant and upon binding undergo a significant alteration of secondary structure. Suitable for practice of the invention are other DSS-like molecules (anionic surfactants like SDS); a wide range of DSS:GLP-1 ratios, for example, 0.1 to 1 to 20:1 or 50:1; a wide range of

-13-

formulation conditions (pH, other non-active excipients, glycerin, alcohol, polymeric additives, coatings, and the like); tablet, liquid or capsule forms; and the like. (Remington's "Pharmaceutical Sciences," 1980).

5

EXAMPLES

The following examples are presented to exemplify, not limit the invention.

Example 1: Preserved Formulations of Val⁸-GLP-1 (7-37) OH with DSS

10 A formulation of the invention was prepared by dissolving Val⁸-GLP-1(7-37)OH at 1 mg/ml in an aqueous solution containing 16 mg/ml glycerin and 10 mM sodium tribasic phosphate. The solution was adjusted to about pH 8.1 using 1N HCl.

15 The preservative m-cresol was prepared at a concentration of 100 mg/ml in absolute ethanol.

Sodium docusate (DSS) was prepared at a concentration of 20 mg/ml in water with gentle warming on a hot plate.

To each of 500 μ L aliquots of the Val⁸-GLP-1(7-37)OH
20 solution in 3-ml glass vials were added 0, 3.3, 6.6 or 16.5 μ L of the DSS solution followed by 15.8 μ L of the preservative m-cresol solution. After gentle mixing of the components in the vials by hand swirling the pH of each clear solution was adjusted to pH 7.8. Replicate samples
25 were incubated at 4°C, ambient temperature, and 37°C. Within 4 hours at ambient temperature, the samples containing 0 or 3.3 μ L of the DSS solution had become hazy due to peptide denaturation.

-14-

After incubation for 16 hours at 37°C, all four types of samples were clear. The solutions were then incubated at 4°C. Again, the solutions containing 0 or 3.3 µL of the DSS solutions became, and remained, hazy.

5 The solutions containing 6.6 µL or 16.5 µL of the DSS solution, which correspond to 2:1 and 10:1 molar ratios DSS to Val⁸-GLP-1(7-37)OH, respectively, remained clear at 4°C for at least 6 weeks. At this time, HPLC analysis showed a purity of the Val⁸-GLP-1(7-37)OH of 98.3% and 97.2%,
10 respectively.

Example 2: A Preserved Formulation of Val⁸-GLP-1 (7-37) OH with DSS

A formulation of the invention was prepared by dissolving Val⁸-GLP-1(7-37)OH at about 1.0 mg/ml in an
15 aqueous solution containing 16 mg/ml glycerin and 10 mM sodium tribasic phosphate. The solution was adjusted to about pH 8.0 using 5N HCl. The solution was then filtered through 0.2 µ and 0.02 µ filters. The peptide concentration was quantified by ultraviolet (UV) analysis at 280 nm.

20 6.5 ml of the Val⁸-GLP-1 solution was added to 1.62 mg of solid DSS, which had been dried from a 100 mg/ml solution in absolute ethanol, to give a 2:1 molar ratio of DSS to Val⁸-GLP-1. After gently stirring 15 minutes at ambient temperature the solution was added to 20.5 mg of m-cresol,
25 which had been dried from a 100 mg/ml solution in absolute ethanol, to give a m-cresol concentration of about 3.15 mg/ml. After stirring 15 minutes at ambient temperature, the solution was adjusted to about pH 7.7 and

-15-

passed through a 0.2 μ filter. Portions of this formulation were stored at 4°C and at ambient temperature.

After 18 weeks, the formulations maintained at 4°C and at ambient temperature were examined. Both solutions were clear. At this time, HPLC analysis showed a purity of the Val⁸-GLP-1(7-37)OH of 98.3% and 90.8% for the 4°C and ambient temperature samples, respectively.

Example 3: In Vivo Effects of a Formulation

A portion of the formulation from Example 2 was injected subcutaneously into beagle dogs that were clamped at an elevated glycemic level (200 mg/dl). 3 nmoles/kg of Val⁸-GLP-1 were injected into each animal. Glucose infusion rates needed to maintain hyperglycemia were measured for 2.5 hours after the injections and compared to injections of a vehicle control solution.

In comparison to the vehicle control, the injection of the Val⁸-GLP-1 formulation resulted in an elevated glucose infusion for about two hours post-injection, indicating appropriate biological activity of the peptide is maintained under these conditions.

Example 4: Preserved Formulations of Val⁸-GLP-1 With Other Surfactants

A formulation of the invention was prepared by dissolving Val⁸-GLP-1(7-37)OH at 1 mg/ml in an aqueous solution containing 16 mg/ml glycerin and 10 mM sodium tribasic phosphate. The solution was adjusted to about pH 8.0 using 2N HCl.

The preservative m-cresol was prepared at a concentration of 100 mg/ml in absolute ethanol.

-16-

Various formulation excipients listed herein were added to 500 μ L aliquots of the Val^s-GLP-1(7-37)OH solution in 3-ml glass vials. After stirring for about 45 minutes at ambient temperature, 15.8 μ l of a 100 mg/ml m-cresol solution in absolute ethanol was added to give a m-cresol concentration of about 3 mg/ml. The test solutions were observed for clarity for about 3 hours at ambient temperature and then at 4°C overnight.

Without any additives, the Val^s-GLP-1 solution becomes hazy at both ambient temperature and at 4°C. Addition of the following surfactants preserved solution clarity at ambient temperature, but not at 4°C: 10 μ l of Tween-40, 10 μ l of Tween-80. Hence these surfactants did improve formulation stability.

-17-

CLAIMS:

1. A formulation comprising a surfactant, and a GLP-1-like peptide.
2. The formulation of Claim 1 wherein the surfactant
5 is selected from the group consisting of DSS (docusate sodium, CAS Registry Number [577-11-7]) and related substances; docusate calcium [CAS number 128-49-4], and docusate potassium [CAS number 7491-09-0].
3. The formulation of Claim 1 wherein the surfactant
10 is selected from the group consisting of SDS (sodium dodecyl sulfate or sodium lauryl sulfate), sodium caprylate, sodium cholate, sodium deoxycholate, sodium taurocholate, and sodium glycocholate.
4. The formulation of Claim 1, wherein the surfactant
15 is selected from the group consisting of zwitterionic (e.g. N-alkyl-N,N-dimethylammonio-1-propanesulfonates, 3-cholamido-1-propyldimethylammonio-1-propane-sulfonate), cationic (cetylpyridinium chloride) non-ionic (Triton[®] X-100, Dodecyl β -D-glucopyranoside), or polymeric
20 (Tween[®]-40, Tween[®]-80, or Brij-35[®]) surfactants.
5. The formulation of Claim 1, wherein the GLP-1-like molecule is selected from the group consisting of
SEQ ID NO:1;
25 SEQ ID NO:4;
SEQ ID NO:5;
peptides of the formula:

-18-

 R_1 -SEQ ID NO:2- R_2

wherein: R_1 is selected from the group consisting of
L-histidine, D-histidine, desamino-histidine,
2-amino-histidine, β -hydroxy-histidine, homohistidine,
5 alpha-fluoromethyl-histidine, and alpha-methyl-histidine,
and R_2 is selected from the group consisting of NH_2 ;

peptides of the formula:

 R_1 -SEQ ID NO:3- R_2

wherein R_1 is selected from the group consisting of
10 4-imidazopropionyl, 4-imidazoacetyl, or 4-imidazo- α , α -
dimethyl-acetyl; R_2 is selected from the group consisting of
Gly-OH or NH_2 . In addition, Lys at position 27 of SEQ ID
NO:3 may be an acyl group selected from the group consisting
of C_6 - C_{10} unbranched acyl or may be absent;
15 and pharmaceutically acceptable salts
thereof.

6. The formulation of Claim 1, further defined as
being stable at a pH of about 6.5 to about 9.0.

7. The formulation of Claim 1, further defined as
20 being stable at about pH 7.0 to about 8.0.

8. The formulation of Claim 1, further comprising an
isotonicity agent.

9. The formulation of Claim 8, wherein the
isotonicity agent is glycerin.

25 10. The formulation of Claim 8, wherein the
isotonicity agent is sodium chloride.

-19-

11. The formulation of Claim 1, further comprising a preservative.

12. The formulation of Claim 11, wherein the preservative is selected from the group consisting of m-
5 cresol, phenol, methylparaben, and benzyl alcohol.

13. A method of treating a person having a condition for which administration of GLP-1 is indicated, said method comprising obtaining a formulation of any one of Claims 1 to 12 and administering a pharmacologically effective amount of
10 the formulation to the person.

14. The method of Claim 13, wherein the condition is diabetes.

15. The method of claim 13, wherein the condition is selected from the group consisting of obesity, myocardial infarction, catabolic states, and stroke.

16. The method of any one of Claims 12 to 13 wherein the administration is oral.

SEQUENCE LISTING

<110> Hoffmann, James A.
Eli Lilly and Company

<120> GLP-1 FORMULATIONS

<130> X-11368

<140>

<141>

<150> US60/067,600

<151> 1997-12-05

<160> 5

<170> PatentIn Ver. 2.0

<210> 1

<211> 31

<212> PRT

<213> Homo sapiens

<400> 1

His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly

1

5

10

15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly

20

25

30

<210> 2

<211> 30

<212> PRT

<213> synthetic construct

<220>

<223> Xaa at position 1 is Ala, Gly, Val, Thr, and Ile;
and Xaa at position 14 is Glu, Gln, Ala, Thr, Ser,
and Gly; and Xaa at position 20 is Glu, Gln, Ala,
Thr, Ser, and Gly;

<220>

<223> and Xaa at position 30 is Gly or absent.

<400> 2

Xaa Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Xaa Gly Gln

1 5 10 15
Ala Ala Lys Xaa Phe Ile Ala Trp Leu Val Lys Gly Arg Xaa
 20 25 30

<210> 3
<211> 30
<212> PRT
<213> synthetic construct

<220>
<223> Xaa at position 19 is Lys or Arg; and Xaa at
 position 30 is Gly or absent; and Lys at position
 27 may be acylated.

<400> 3
Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly Gln
1 5 10 15

Ala Ala Xaa Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Xaa
 20 25 30

<210> 4
<211> 30
<212> PRT
<213> Homo sapiens

<220>
<223> Arg at position 30 is C-terminally amidated.

<400> 4
His Ala Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg
 20 25 30

<210> 5
<211> 31
<212> PRT
<213> synthetic construct

<400> 5
His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
20 25 30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/25515

A. CLASSIFICATION OF SUBJECT MATTER		
IPC(6) :A61K 38/26 US CL :514/12, 21 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) U.S. : 514/12, 21, 530/308, 324		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, DIALOG, DERWENT DWPI, STN search terms: glucagon like peptide, glp, oral, surfactant, docusate, SEQ ID NO:2		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	WO 93/18785 A1 (NOVO NORDISK A/S) 30 September 1993, page 2, lines 17-21, page 5, line 12 -page 6, line 31, page 8, lines 25-28, page 10, lines 1-26.	1, 2, 4-14 --- 15
Y	WO 97/31943 A1 (NOVO NORDISK A/S) 04 September 1997, page 3, lines 22-25.	15
X	US 5,120,712 A (HABENER) 09 June 1992, column 7, lines 3-6, claim 8.	1, 2, 5, 13, 14
X	US 5,376,637 A (SAWAI ET AL) 27 December 1994, column 1, lines 19-20, column 3, lines 28-31, column 4, lines 26-28, Medicine Preparation Examples 1, 5, 8.	1-4, 6-8, 10, 11, 13, 16
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
A	document defining the general state of the art which is not considered to be of particular relevance	*T*
E	earlier document published on or after the international filing date	*X*
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*
O	document referring to an oral disclosure, use, exhibition or other means	
P	document published prior to the international filing date but later than the priority date claimed	*A*
	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
	document member of the same patent family	
Date of the actual completion of the international search 26 JANUARY 1999		Date of mailing of the international search report 11 MAR 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer JEFFREY E. RUSSEL Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/25515

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,545,618 A (BUCKLEY ET AL) 13 August 1996.	1-16
X,P	US 5,766,620 A (HEIBER ET AL) 16 June 1998, column 5, lines 62-67, column 6, lines 1-7, 30-31, 60-63, column 7, lines 28-45, column 13, lines 29-61.	1-3, 5-7, 13, 14, 16
---		---
Y,P		15
X,P	US 5,811,388 A (FRIEND ET AL) 22 September 1998, Abstract, column 9, lines 7-22, column 9, line 61 - column 10, line 21.	1, 2, 6, 7, 13, 16
---		---
Y,P		15